

Walmart Inc.


-PE Product Labels –

-continued











3821









DAYTIME SEVERE

Cold & Flu

Acetaminophen - Pain Reliever/Fever Reducer
Dextromethorphan HBr - Cough Suppressant
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

MAXIMUM STRENGTH

Relief of:

- Headache, fever, sore throat, minor aches and pains
- Nasal/sinus congestion and sinus pressure
- Chest congestion
- Cough

Actual Size

24 SOFTGELS

MAXIMUM STRENGTH

Cold & Flu

DAYTIME SEVERE

equate

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg, Pain reliever/fever reducer
Dextromethorphan HBr 10 mg, Cough suppressant
Guaifenesin 200 mg, Expectorant
Phenylephrine HCl 5 mg, Nasal decongestant

Purpose

Use ■ temporarily relieves common cold/symptoms ■ nasal congestion ■ sinus congestion & pressure ■ cough due to minor throat & bronchial irritation ■ minor aches & pains ■ headache ■ fever ■ sore throat ■ reduces swelling of nasal passages ■ temporarily restores free breathing through the nose ■ promotes nasal/sinus drainage ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both excess mucus and make coughs more productive.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 softgels in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ If you are now taking a prescription antidepressant called MAOI (certain drugs for depression, psychiatric or emotional conditions, or

Drug Facts (continued)

Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ take only as directed

■ do not exceed 8 softgels per 24 hours

adults & children 12 years & over	2 softgels with water every 4 hours
children 4 to under 12 years	ask a doctor
children under 4 years	do not use

Other information ■ do not exceed 25°C (77°F)

Inert ingredients FD&C Yellow #6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-387-1915.

DISTRIBUTED BY:
Wal-Mart Inc.,
 Bentonville, AR 72716
PRODUCT OF INDIA
 This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademarks **Vicksal** DayQuil® Severe Cold & Flu.

Compare to Vicksal DayQuil® Severe Cold & Flu active ingredients*

NEC7985-162-24


24 SOFTGELS

Drug Facts (continued)
Questions or comments?
1-888-387-1915

LOT NO:
EXP DATE:

6 81313 10063 2


TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

NO WATERMARK ON PRINTING
 LOT # & EXP. DATE AREA
 ☆ 729565

 6 81131 37878 9

Maximum Strength

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
 *This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucine® FAST-MAX® Day Time Cold & Flu and Night Time Cold & Flu. 50844 ORG071961769401
 PARENTS: www.StarMedicationHouse.org

Daytime


 ND 49035-358-01

Compare to Maximum Strength Mucine® FAST-MAX® Day Time Cold & Flu and Night Time Cold & Flu Active Ingredients*

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

Nighttime

Daytime

DAYTIME
SEVERE
MULTI-SYMPTOM

Cold

Acetaminophen -
Pain Reliever/Fever Reducer
Dextromethorphan HBr -
Cough Suppressant
Phenylephrine HCl -
Nasal Decongestant

NIGHTTIME
SEVERE
Cough

Acetaminophen -
Pain Reliever/Fever Reducer
Diphenhydramine HCl -
Antihistamine/Cough Suppressant
Phenylephrine HCl -
Nasal Decongestant

DAYTIME & NIGHTTIME RELIEF VALUE PACK

DAYTIME
SEVERE
MULTI-SYMPTOM

Cold

Acetaminophen -
Pain Reliever/Fever Reducer
Dextromethorphan HBr -
Cough Suppressant
Phenylephrine HCl -
Nasal Decongestant

Dye free • Aspartame free • Sodium free

Temporarily Relieves

- Nasal congestion • Body aches
- Sore throat/pain • Fever
- Cough • Headache

Green tea & honey lemon flavors

Infused with menthol & green tea flavors

Naturally and Artificially Flavored

6 PACKETS

NIGHTTIME
SEVERE

Cold & Cough

Acetaminophen -
Pain Reliever/Fever Reducer
Diphenhydramine HCl -
Antihistamine/Cough Suppressant
Phenylephrine HCl -
Nasal Decongestant

Aspartame free • Sodium free

Temporarily Relieves

- Nasal congestion • Body aches
- Sore throat/pain • Fever
- Cough • Runny nose
- Headache • Sneezing

Honey lemon flavor

Infused with chamomile & white tea flavors

Naturally and Artificially Flavored

6 PACKETS

12 TOTAL PACKETS

Lot: _____

Exp: _____

DATE/TIME: _____

TAMPER EVIDENT INNER UNIT: DO NOT USE IF SEALED PACKET IS TORN OR BROKEN.

© 2017

6 81131 26822 4


PRODUCED BY: Belmont Inc.
Burlington, NJ 07718

Pharmaceutical Research and Development of
Pharmaceuticals and the manufacture of
Pharmaceuticals are trademarks of Belmont Inc.
© 2017 Belmont Inc. All rights reserved.
Pharmaceutical Research and Development of
Pharmaceuticals are trademarks of Belmont Inc.

RECYCLE
PAKES
DO NOT
RECYCLE

MADE IN THE USA

3327



Compare to Tylenol® Allergy Multi-Symptom Active Ingredients*

Rapid Release

Multi-Symptom Allergy Relief

Pain reliever, Nasal decongestant, Antihistamine

Drug Facts (continued)

Uses ■ temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ sinus congestion and pressure ■ nasal congestion ■ runny nose and sneezing ■ headache ■ minor aches and pains

Drug Facts (continued)

When using this product ■ do not exceed recommended dosage ■ excitability may occur, especially in children ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ drowsiness may occur

Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain or nasal congestion gets worse or lasts more than 7 days ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning) ■ adults and children 12 years and over ■ take 2 gelscaps every 4 hours ■ do not take more than 12 gelscaps in 24 hours ■ children under 12 years: do not use this adult product in children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage

Other information ■ see end flap for expiration date and lot number

Drug Facts

Active ingredients (in each gelscap) Purpose

Acetaminophen 325 mg. Pain reliever

Chlorpheniramine maleate 2 mg. Antihistamine

Phenylephrine HCl 5 mg. Nasal decongestant

Drug Facts (continued)

Warnings ■ temporarily relieves these additional symptoms of hay fever: ■ itching of the nose or throat ■ itchy, watery eyes ■ helps clear nasal passages ■ helps decongest sinus openings and passages

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 12 gelscaps in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have ■ liver disease ■ diabetes ■ heart disease ■ a breathing problem such as emphysema or chronic bronchitis ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Drug Facts (continued)

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ contains FD&C Yellow #5 (tartrazine) as a color additive

Inactive ingredients croscarmellose sodium, additive

Drug Facts (continued)

croscarmellose, FD&C red #3, FD&C red #40, FD&C yellow #5, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol.

Drug Facts (continued)

shellic glue, silica gel, stearic acid, titanium dioxide

Questions? 1-888-287-1915

Drug Facts (continued)

Other information ■ see end flap for expiration date and lot number

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Ask a doctor before use if you have ■ liver disease ■ diabetes ■ heart disease ■ a breathing problem such as emphysema or chronic bronchitis ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Drug Facts (continued)

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ contains FD&C Yellow #5 (tartrazine) as a color additive

Inactive ingredients croscarmellose sodium, additive

Drug Facts (continued)

croscarmellose, FD&C red #3, FD&C red #40, FD&C yellow #5, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol.

Drug Facts (continued)

Other information ■ see end flap for expiration date and lot number

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Ask a doctor before use if you have ■ liver disease ■ diabetes ■ heart disease ■ a breathing problem such as emphysema or chronic bronchitis ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

Drug Facts (continued)

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ contains FD&C Yellow #5 (tartrazine) as a color additive

Inactive ingredients croscarmellose sodium, additive


Drug Facts (continued)

croscarmellose, FD&C red #3, FD&C red #40, FD&C yellow #5, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol.

Satisfaction guaranteed - for questions or comments please call 1-888-287-1915.
 Distributed by: Wal-Mart Stores, Inc., Bentonville, AR 72716
 © Wal-Mart Stores, Inc.
 *This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® Allergy Multi-Symptom.
 50844 REV0812455908
 6 81131 03925 3
 CN 2014 0055

B-2203-559-08
 REV0812455908

08-655-5205-00



Maximum Strength

Severe Allergy Plus Sinus Headache

**Acetaminophen, Diphenhydramine HCl,
Phenylephrine HCl**



Equate 44-543

[illegible]

NDC 49035-165-09

Compare to Maximum Strength Mucinex® SINUS-MAX® Severe Congestion & Pain Active Ingredients*

Maximum Strength Sinus Severe Congestion Relief

Acetaminophen - Pain Reliever
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

- Relieves headache
- Thins & loosens mucus
- Relieves sinus congestion

20 CAPLETS

For Ages 12+

Actual Size

Drug Facts (continued)

Other information ■ each caplet contains sodium 3 mg

Directions ■ do not take more than directed

■ children under 12 years: do not use

■ adults and children 12 years and over: take 2 caplets every 4 hours

■ do not take more than 12 caplets in any 24-hour period

■ signs or symptoms:

critical for adults as well as for children even if you do not notice any

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is

If pregnant or breast-feeding, ask a health professional before use.

could be signs of a serious condition.

■ cough comes back or occurs with rash or headache that lasts. These

■ new symptoms occur

■ redness or swelling is present

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ nervousness, dizziness, or sleepiness occur

Stop use and ask a doctor if

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg.....Pain reliever

Guaifenesin 200 mg.....Expectorant

Phenylephrine HCl 5 mg.....Nasal decongestant

Purpose

Uses ■ temporarily relieves: ■ nasal congestion ■ headache

■ sinus congestion and pressure ■ minor aches and pains

■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the

■ temporarily promotes nasal and/or sinus drainage

Drug Facts (continued)

alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-888-287-1915

Drug Facts (continued)

Warnings

This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ liver disease ■ thyroid disease

■ heart disease ■ diabetes

■ high blood pressure

■ difficulty in urination due to enlargement of the prostate gland

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Inactive ingredients corn starch, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl

Drug Facts (continued)

Other information ■ each caplet contains sodium 3 mg

Directions ■ do not take more than directed

■ children under 12 years: do not use

■ adults and children 12 years and over: take 2 caplets every 4 hours

■ do not take more than 12 caplets in any 24-hour period

■ signs or symptoms:

critical for adults as well as for children even if you do not notice any

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is

If pregnant or breast-feeding, ask a health professional before use.

could be signs of a serious condition.

■ cough comes back or occurs with rash or headache that lasts. These

■ new symptoms occur

■ redness or swelling is present

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ nervousness, dizziness, or sleepiness occur

Stop use and ask a doctor if

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Other information ■ each caplet contains sodium 3 mg

Directions ■ do not take more than directed

■ children under 12 years: do not use

■ adults and children 12 years and over: take 2 caplets every 4 hours

■ do not take more than 12 caplets in any 24-hour period

■ signs or symptoms:

critical for adults as well as for children even if you do not notice any

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is

If pregnant or breast-feeding, ask a health professional before use.

could be signs of a serious condition.

■ cough comes back or occurs with rash or headache that lasts. These

■ new symptoms occur

■ redness or swelling is present

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ nervousness, dizziness, or sleepiness occur

Stop use and ask a doctor if

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Warnings

This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ liver disease ■ thyroid disease

■ heart disease ■ diabetes

■ high blood pressure

■ difficulty in urination due to enlargement of the prostate gland

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Inactive ingredients corn starch, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl

Drug Facts (continued)

Other information ■ each caplet contains sodium 3 mg

Directions ■ do not take more than directed

■ children under 12 years: do not use

■ adults and children 12 years and over: take 2 caplets every 4 hours

■ do not take more than 12 caplets in any 24-hour period

■ signs or symptoms:

critical for adults as well as for children even if you do not notice any

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is

If pregnant or breast-feeding, ask a health professional before use.

could be signs of a serious condition.

■ cough comes back or occurs with rash or headache that lasts. These

■ new symptoms occur

■ redness or swelling is present

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ nervousness, dizziness, or sleepiness occur

Stop use and ask a doctor if

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

Drug Facts (continued)

Warnings

This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

■ liver disease ■ thyroid disease

■ heart disease ■ diabetes

■ high blood pressure

■ difficulty in urination due to enlargement of the prostate gland

■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

■ cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Inactive ingredients corn starch, croscopolone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING


No print/No varnish area
Lot no/Exp date


B-2203-615-09-SM
REV0623861509

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915.

DISTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Maximum Strength Mucinex® SINUS-MAX® Severe Congestion & Pain. 50344 REV0623861509





KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts (continued)

Active ingredients (in each gelcap)

Acetaminophen 325 mg. Pain reliever/fever reducer
 Pseudoephedrine HCl 5 mg. Nasal decongestant

Purpose

Temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold: headache, nasal congestion, sinus congestion and pressure, minor aches and pains, helps decongest sinus openings and passages, promotes sinus drainage, helps clear nasal passages, temporarily reduces fever.

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ rash ■ skin redness ■ blisters ■ If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Directions

do not take more than directed ■ adults and children 12 years and over ■ take 2 gelcaps every 4 hours ■ do not take more than 10 gelcaps in 24 hours ■ children under 12 years: ask a doctor

Other information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Questions or comments? 1-888-287-1915

Inactive ingredients

croscarmellose sodium, croscopolone, D&C yellow #6, #10, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropylcellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyethylene glycol, shellac, silica, silicon dioxide, stearic acid, titanium dioxide

Questions or comments? 1-888-287-1915

Drug Facts (continued)

■ avoid high humidity ■ see end flap for expiration date and lot number

Lot/Exp area at minimum size 1.25 x .5"

729655
 6 81131 29795 0

DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

TAMPER EVIDENT:

DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Satisfaction guaranteed - Or we'll replace it or give you your money back.

For questions or comments, please call 1-888-287-1915.

DISTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716

owner of the registered trademark Tylenol®

SINUS + HEADACHE Day.

50844 REV0418053822

3336

B-2203-558-22-R
REV 04/18/05/8822**equate™**

Rapid Release

**Sinus
Congestion & Pain**

NDC 45015-558-22

equate™

Rapid Release

**Sinus
Congestion
& Pain****Acetaminophen, Phenylephrine HCl**

Pain Reliever, Fever Reducer, Nasal Decongestant

Daytime Non-Drowsy

- Sinus headache & pressure
- Nasal congestion

**Compare to
Tylenol® SINUS
+ HEADACHE
Day Active
Ingredients*****48**

GELCAPS

Actual Size



NO. 4503-527-08

DAYTIME

Sinus Congestion & Pain

Acetaminophen - Pain Reliever/Fever Reducer
Guaifenesin - Expectorant
Phenylephrine HCl - Nasal Decongestant

Relieves:

- Sinus headache
- Sinus pressure
- Nasal congestion
- Mucus & chest congestion

24 CAPLETS

Drug Facts
Active ingredients (in each caplet)
 Acetaminophen 325 mg... Pain reliever/fever reducer
 Guaifenesin 200 mg... Expectorant
 Phenylephrine HCl 5 mg... Nasal decongestant

Purpose
 Relieves sinus headache, sinus pressure, nasal congestion, mucus & chest congestion, and pain.

Directions
 Take 2 caplets every 4 hours.

Warnings
 Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Other information
 ■ **TAMPER EVIDENT, DO NOT USE IF OUTER PACKAGE IS OPENED OR BUSTEN IS TORN OR BROKEN**

Drug Facts (continued)

■ See and keep for expiration date and lot number

Inactive ingredients corn starch, croscarmellose, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol.

Questions or comments? 1-888-287-1915

Drug Facts (continued)

polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

No Print/No Varnish Lot and Expiration No.

Satisfaction guaranteed - For questions or comments please call 1-888-287-1915

TAMPER EVIDENT, DO NOT USE IF PACKAGE IS OPENED OR IF BUSTEN UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DISTRIB BY: Oxy-Pharm, Inc., Boston, MA 02118
 © 2008 Oxy-Pharm, Inc. All rights reserved.
 Oxy-Pharm, Inc. is a registered trademark of Oxy-Pharm, Inc.

Drug Facts (continued)

■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ hives ■ rash ■ skin redness

If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

■ If you have ever had an allergic reaction to this product or any of its ingredients

B-2203-527C-08-SC
 REV0922M52708